



CORRECTION

Open Access



Correction to: Comparison of the effects of olfactory stimulation by air-dried and high-temperature-dried wood chips of hinoki cypress (*Chamaecyparis obtusa*) on prefrontal cortex activity

Harumi Ikei^{1,2}, Chorong Song¹, Juyoung Lee³ and Yoshifumi Miyazaki^{1*}

Correction to: *J Wood Sci* (2015) 61:537–540

<https://doi.org/10.1007/s10086-015-1495-6>

After the article was published [1], an error was found in Fig. 2. The following change should be made to the figure in the published article. The change does not affect the conclusions of the article in any way.

Figure 2 should be replaced with the following figure:




Fig. 2 Time-dependent oxy-Hb concentration changes (per second) in the prefrontal cortex during olfactory stimulation by air-dried or high-temperature-dried hinoki cypress wood chips. The oxy-Hb concentration shown is the difference between the pre- and postmeasurement conditions. Data are expressed as mean \pm SE; $n = 19$

*Correspondence: ymiyazaki@faculty.chiba-u.jp

¹ Center for Environment, Health, and Field Sciences, Chiba University, 6-2-1 Kashiwa-no-ha, Kashiwa, Chiba 277-0882, Japan

Full list of author information is available at the end of the article

Author details

¹ Center for Environment, Health, and Field Sciences, Chiba University, 6-2-1 Kashiwa-no-ha, Kashiwa, Chiba 277-0882, Japan. ² Present Address: Forestry and Forest Products Research Institute, 1 Matsunosato, Tsukuba, Ibaraki 305-8687, Japan. ³ Korea Forest Service Republic of Korea, Government Complex-Daejeon Bldg. 1, 189 Cheongsa-ro, Seo-gu, Daejeon 302-701, Republic of Korea.

The original article can be found online at <https://doi.org/10.1007/s10086-015-1495-6>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Published online: 09 March 2019

Reference

1. Ikei H, Song C, Lee J, Miyazaki Y (2015) Comparison of the effects of olfactory stimulation by air-dried and high temperature-dried wood chips of hinoki cypress (*Chamaecyparis obtusa*) on prefrontal cortex activity. *J Wood Sci* 61:537–540. <https://doi.org/10.1007/s10086-015-1495-6>